arXiv:1510.06877v1 [astro-ph.SR] 23 Oct 2015

Evidence of Resonant Mode Coupling in the Pulsating DB White Dwarf Star KIC 08626021

W. Zong[1], S. Charpinet[1] and G. Vauclair[1]

[1] *CNRS, Université de Toulouse, UPS-OMP, IRAP, Toulouse 31400, France;*
weikai.zong@irap.omp.eu , stephane.charpinet@irap.omp.eu ,
gerard.vauclair@irap.omp.eu

Abstract. The *Kepler* spacecraft provides new opportunities to search for long term frequency and amplitude modulations of oscillation modes in pulsating stars. We analyzed nearly two years of uninterrupted data obtained with this instrument on the DBV star KIC 08626021 and found clear signatures of nonlinear resonant mode coupling affecting several triplets. The behavior and timescales of these amplitude and frequency modulations show strong similarities with theoretical expectations. This may pave the way to new asteroseismic diagnostics, providing in particular ways to measure for the first time linear growth rates of pulsation modes in white dwarf stars.

1. Introduction

The *Kepler* spacecraft is a magnificent instrument to search for long term frequency and amplitude modulations of oscillation modes in pulsating stars. Among the 6 pulsating white dwarfs present in the *Kepler* field, KIC 08626021 is the unique DB pulsator. It has a rotation period $P_{rot} \sim 1.7$ days, estimated from the observed frequency spacings of 3 g-mode triplets (Østensen et al. 2011). It has been observed by *Kepler* for 23 months in short cadence (SC) mode without interruption. Thus, it is a suitable candidate to investigate the resonant mode coupling mechanisms that could induce long term amplitude and frequency modulations of the oscillation modes. Such resonant couplings are predicted to occur in triplets where the rotationally shifted components have frequencies ν_1 and ν_2 such that $\nu_1 + \nu_2 \sim 2\,\nu_0$, where ν_0 is the frequency of the central component. The theoretical exploration of those mechanisms was extensively developed long before the era of space observations (Buchler et al. 1995, 1997) but was almost interrupted more than a decade ago because of the lack of clear observational evidence of such phenomena, due to the difficulty of capturing amplitude or frequency variations that occur on months to years timescales from ground based observatories. Resonant coupling within triplets was proposed for the first time as the explanation for the frequency and amplitude long term variations observed in the GW Vir pulsator PG 0122+200 (Vauclair et al. 2011). We present the analysis of KIC 08626021 in which two triplets exhibit amplitude and frequency variations during the 23-month of observation. Such time modulations pave the way to new asteroseismic diagnostics, providing in particular ways to measure for the first time linear growth rates of pulsation modes in white dwarf stars.

2. Frequency and amplitude modulations

The white dwarf star KIC 08626021 has been continuously observed by *Kepler* since quarter Q10.1 up to Q17.2. The high precision photometric data cover ~ 684.2 days (23 months), with a duty circle of ~ 89 %. We used a dedicated software, FELIX, to extract frequencies (details of the program can be found in Charpinet et al. 2010). In this study, we concentrate on rotationally split triplets and investigate the variation of amplitude and frequency between components of these triplets and their relationship. We point out that the frequencies near 3682 μHz reported by Østensen et al. (2011) were in fact resolved into several close peaks with the 23-month light curve. It is therefore probably not a real triplet contrary to the other structures found at 4310 and 5073 μHz (see below). In order to study the variability with time of these modes, we constructed a filter window covering 200 days and slid the filter window along the whole light curve by time steps of 20 days, thus constructing a time-frequency diagram. We also prewhithened the frequencies "chunk by chunk", i.e., the 23-month light curve of KIC 08626021 was divided into 20 chunks, each containing 6-month of data except the last 3 chunks being at the end of the observations. The results for the two triplets are discussed below.



Figure 1. Frequency and amplitude modulations of the triplet at 4310 μHz. The frequencies and amplitudes of the triplet show clear signatures of periodic modulations, as shown in the left bottom panel and right panel. The grey scale in the upper left panel represents the amplitude. The dashed line in the lower right panel is the average value of the frequency (see text for details). Note that the frequency scales (x axis) in the upper and lower left panels are different.

The amplitude and frequency modulations of the triplet near 4310 μHz are shown in Figure 1 . The grey scale (or color scale for electronic version) in the upper left panel represents the amplitude. The vertical dashed line in the lower left panel is the average


T-F Periodogram (prewhitened)
Amplitude (%) vs Frequency (uHz) & Time (d)
Time (BJD-2455000 d)
Frequency (μHz)
5070.0307 μHz (m = -1)
5073.234 μHz (m = 0)
5076.4354 μHz (m = +1)
Amplitude (%)
Time (BJD-2455000 d)

Figure 2. Frequency and amplitude modulations as shown in Figure 1 but for the triplet at 5073 μHz.The amplitudes of the triplet show clear signatures of periodic modulations, as shown in the right panel. The frequencies of the triplet are relatively stable during the observation.

value of the frequency over the entire run. The right panel shows the amplitude modulations of each component forming the triplet. Both the amplitudes and frequencies show clear signatures of quasi periodic modulations with the same timescale of ~ 750 days. The frequencies and amplitudes of the side components evolve in phase and are antiphased with the central component. Figure 2 shows the modulations observed in the other triplet at 5073 μHz. The frequencies in this triplet appear to be stable during the nearly two years of monitoring, while the amplitudes show modulations. Note that the amplitude of the m = -1 component went down below the 4σ detection threshold and was essentially lost in the noise during the last half of the observations.

3. Discussion

The frequency and amplitude modulations observed in the triplets of KIC 086226021 can be related to nonlinear resonant mode coupling mechanisms. The first triplet at 4310 μHz (Figure 1) behaves like if it is in the intermediate regime of the resonance, in which the oscillation modes undergo periodic amplitude and frequency modulations. Theory suggests that the time scale of these modulations should be roughly a few times the inverse of the growth rate of the pulsating mode (Goupil et al. 1998). Therefore this periodicity could in principle be used to measure the growth rate. In addition if we compare the second order effect of rotational splitting as estimated directly from the measured mean frequencies :

$$\delta\nu = \nu_1 + \nu_2 - 2\nu_0 \qquad (1)$$

with the estimated value following Dziembowski & Goode (1992),

$$\delta\nu = 4C\frac{\Omega^2}{\nu_0} \qquad (2)$$

where C is the first order Ledoux constant (~ 0.5 for dipole g-modes) and $\Omega = 2\pi/P_{rot}$ is the angular frequency of the stellar rotation, i.e ., $\delta\nu \sim 0.018\,\mu$Hz, both are found to be very similar. In the intermediate regime, the expected periodic modulation timescale is $P_{mod} \sim 1/\delta\nu$ (Goupil et al. 1998), which with the values given above leads to $P_{mod} \sim$ 650 days, i.e., very similar to the amplitude and frequency modulation timescale of 750 days roughly estimated from Figure 1. This further supports our interpretation that nonlinear resonant coupling is indeed at work in this star. The triplet at $5073\,\mu$Hz (Figure 2) would be in a different regime, likely in the nonresonant regime. The amplitude of the components show clear modulations while the frequency are relatively stable during the observation. This means the ratio of the real part over the imaginary part of the coupling coefficients is large in that case. This ratio roughly measures nonlinear nonadiabaticities in the star. Hence our result shows that two neighbor triplets can belong to different resonant regimes (frequency lock, time dependent or nonresonant), as it was also suggested in the white dwarf star GD 358 (Goupil et al. 1998).

4. Conclusion

Frequency and amplitude modulations of oscillation modes have been found in several rotationally split multiplets detected in the DB pulsator KIC 08626021, thanks to the high quality and long duration photometric data obtained with the *Kepler* spacecraft. These modulations show signatures pointing toward nonlinear resonant coupling mechanisms occuring among the multiplet components. This is the first time that such signatures are identified so clearly in white dwarf pulsating stars. Periodic modulations of frequency and amplitude that occur in the intermediate resonant regime may allow for new asteroseismic diagnostics, providing in particular a way to measure for the first time linear growth rates of pulsation modes in white dwarf stars. Such results should motivate further theoretical work on nonlinear resonant mode coupling mechanisms and revive interest in nonlinear stellar pulsation theory in general. Finally, we mention that similar modulations are also found in hot B subdwarf stars according to *Kepler* data.

Acknowledgments. WKZ acknowledges the financial support from the China Scholarship Council. This work was supported in part by the Programme National de Physique Stellaire (PNPS, CNRS/INSU, France) and the Centre National d'Etudes Spatiales (CNES, France).

References

Buchler, J. R., Goupil, M.-J. & Hansen, C. J. 1997, A&A, 321, 159
Buchler, J. R., Goupil, M.-J. & Serre, T. 1995, A&A, 296, 405
Charpinet, S., Green, E. M., Baglin, A., et al. 2010, A&A, 516, L6
Dziembowski, W. A. & Goode, Philip R. 1992, ApJ, 394, 670
Goupil, M. J., Dziembowski, W. A. & Fontaine, G. 1998, BaltA, 7, 21
Østensen, R. H., Bloemen, S., Vučković, M., et al. 2011, ApJL, L39, 736
Vauclair, G., Fu, J.-N., Solheim, J.-E., et al. 2011, A&A, 528, 5